

# Procedure note for Neonatal Circumcision

**Preoperative diagnosis:** Desires Neonatal Circumcision

**Postoperative diagnosis:** same

**Procedure:** Neonatal Circumcision

**Surgeon(s):** \_\_\_\_\_

**Preprocedure counseling:** The risks, benefits, and alternatives of the procedure were discussed with the patient's parent/guardian.

**Procedure:**

A timeout was performed prior to starting the procedure.

The infant was laid in a supine position and the surgical field was prepped and draped in usual sterile fashion. A pacifier with sucrose water was used to aid anesthesia.

\_\_\_\_\_ mL of 1% lidocaine without epinephrine was used to anesthetize the penis with a

**dorsal penile nerve block      subcutaneous ring block      (circle one)**

A dorsal slit was made after clamping the foreskin. The foreskin was retracted and adhesions were removed bluntly. The

**1.1    1.3    1.6 cm      (circle one)**

Gomco clamp was placed in usual fashion ensuring the dorsal slit was completely included and that the amount of foreskin was symmetric on all sides. After securing the Gomco clamp to ensure hemostasis, the foreskin was cut with a scalpel. The Gomco clamp was removed. Hemostasis was assured. The wound was dressed with 1/2" petrolatum gauze.

The attending physician **Dr.** \_\_\_\_\_ was present throughout the entire procedure.

Additional comments:

Signature: \_\_\_\_\_ ID#: \_\_\_\_\_ Date: \_\_\_\_\_ Time: \_\_\_\_\_



**VENTURA COUNTY MEDICAL CENTER  
SANTA PAULA HOSPITAL  
CIRCUMCISION PROCEDURE NOTE**

Addressograph (Two patient identifiers required):